



GEOLOGICAL CO₂ STORAGE

Storing carbon dioxide (CO₂) underground is not a new or emerging technology—it is an existing reality on an industrial scale. In fact, there are geological systems that naturally contain CO₂ and many others throughout the world that experts have determined can retain centuries' worth of injected CO₂. This will help abate climate change by removing and keeping this greenhouse gas out of the atmosphere.

HOW DOES GEOLOGICAL STORAGE OF CO₂ WORK?


Geological storage involves injecting CO₂ captured from industrial processes into rock formations deep underground. The formations are selected for their huge capacity to store and retain the injected greenhouse gases indefinitely. This way the CO₂ is effectively removed and isolated from the atmosphere.

The following geologic characteristics are associated with effective storage sites:

- storage formations have enough voids, or pores, in the rock to allow the injection of CO₂
- pores in the rock are connected well enough, a feature called 'permeability', so that the CO₂ can move and spread out within the formation, providing the capacity to accept the needed amount of CO₂
- formations must have an extensive cap or barrier at the top to contain the CO₂ for hundreds to thousands of years, and longer.

Fortunately, there are many locations around the world that have formations with these characteristics, most of which are found in vast geological features called 'sedimentary basins'. Almost all oil and gas production is associated with sedimentary basins; the types of geologic formations

that trap oil and gas (and also naturally occurring CO₂) include sandstones, limestones and dolomites that are similar to those that make good CO₂ storage reservoirs. It is the natural geologic characteristics that resulted in oil and gas being trapped for millions of years before they were discovered that make secure geologic storage of CO₂ such a viable option for greenhouse gas mitigation.



HOW IS CO₂ INJECTED UNDERGROUND AND WHY DOES IT STAY THERE?

Once captured, the CO₂ is compressed into a fluid almost as dense as water, known as a 'supercritical state', and then pumped down through a well into a porous geological formation as described above. The pores in underground formations are initially filled with a fluid, either oil, gas or, much more commonly, very salty water. Although CO₂ can be injected into oil reservoirs to help with oil recovery, most future large-scale CO₂ injection projects will target a saline water-bearing formation for storage because they are more common and can have enormous capacity. In general, depths greater than 800 metres are desired to keep the CO₂ in the compressed, or dense, state.

Because the CO₂ is initially slightly more buoyant than water, a portion will migrate to the top of the formation, where it will become trapped beneath the caprock, which acts as a seal. In most natural systems, there are numerous thick barriers between the deep reservoir and the surface. Some of the CO₂ will start to dissolve slowly into the saline water and become effectively trapped indefinitely, whereas another portion of CO₂ may become residually trapped in tiny pore spaces. The ultimate trapping process involves dissolved CO₂ reacting with the reservoir rocks to form a mineral, much like snails or clams use calcium and carbon from seawater to form their hard shells. Depending on the reservoir minerals present, this process can be relatively quick or very slow, but it effectively transforms the CO₂ into a solid mineral.

IS UNDERGROUND STORAGE OF CO₂ SAFE?

Three industrial-scale storage projects injecting up to two million tonnes of CO₂ annually into saline formations have been operating for many years, along with other smaller projects actively capturing and storing CO₂.

These industrial-level experiences are complemented by numerous research-scale CCS projects, intergovernmental and industry partnerships, research programs, and stakeholder networks. No adverse safety, health, or environmental effects have ever been documented from any of these operations.

HOW DO WE KNOW THAT IT WORKS?

There are decades of operational experience from projects that are very similar to CCS, including underground CO₂ injection for enhanced oil recovery (EOR) and the use of technologies analogous to CCS, such as acid gas (a combination of hydrogen sulphide and CO₂) injection, and natural gas storage.

The oil and natural gas industries have more than 40 years' experience injecting CO₂ into geologic reservoirs to increase

oil production. This process is a type of EOR and uses the properties of CO₂ to mix with the oil to move it out of the reservoir more effectively. In most operations, the CO₂ is recycled and will remain in the reservoir indefinitely at the end of the life of the oil field. These sites have been injecting many millions of tonnes of CO₂ safely into the subsurface for decades.

The success of these projects and the increasing number of research demonstrations provides considerable confidence in the potential to store large quantities of CO₂ underground safely, securely and for very long periods.

HOW MUCH CO₂ CAN BE STORED UNDERGROUND?

The United Nations Intergovernmental Panel on Climate Change estimates the world's potential capacity at two trillion tonnes, although there could be a "much larger potential".¹

Several regions around the world—the United States, Canada, China, South Africa, Europe and Australia—are doing significant amounts of work on characterising potential storage sites.

The 2012 North American Carbon Storage Atlas indicates there is more than 1600 billion tonnes of CO₂ storage potential in saline formations alone in the United States, Canada and Mexico. This means that there is centuries' worth of CO₂ geological storage for the region.

A European Union project estimates the CO₂ storage capacity in oil and gas fields, in and around the North Sea alone, at 37 billion tonnes, which would enable this region to inject CO₂ for several decades once the fields are depleted.²

The Sleipner project, located some 240 kilometres off the coast of Norway in the North Sea, is storing more than 2,700 tonnes of CO₂ per day, injected nearly 800 metres below the seabed. Over the lifetime of the project, it is expected that more than 20 million tonnes of CO₂ will be injected into the saline formation.

Monitoring surveys of the injected CO₂ indicate that over the past 15 years, the gas has spread out over nearly 10 square kilometres underground, without moving upwards or out of the storage reservoir.

Long-term simulations also suggest that over hundreds of thousands of years, the CO₂ will eventually dissolve in the saline water, becoming heavier and less likely to migrate away from the reservoir.

1 U.N. IPCC, 'Special Report on Carbon Dioxide Capture and Storage: Summary for Policymakers,' 2005, 12.

2 European Union Fifth Framework Programme for Research and Development, 'Geological Storage of CO₂ from Combustion of Fossil Fuels,' November 2004, 9.

FOR MORE INFORMATION

GLOBALCCSINSTITUTE.COM OR EMAIL INFO@GLOBALCCSINSTITUTE.COM